

兔抗 TNFSF12 多克隆抗体

中文名称: 兔抗 TNFSF12 多克隆抗体

英文名称: Anti-TNFSF12 rabbit polyclonal antibody

别 名: APO3L; DR3LG; TWEAK

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

抗原: TNFSF12

宿 主: Rabbit

反应种属: **Human Mouse**

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	The protein encoded by this gene is a cytokine th at belongs to the tumor necrosis factor (TNF) liga nd family. This protein is a ligand for the FN14/T WEAKR receptor. This cytokine has overlapping sig naling functions with TNF, but displays a much wi der tissue distribution. This cytokine, which exists in both membrane-bound and secreted forms, can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells, and thus acts as a regulator of angiogenesis.
Applications:	WB, IHC, IF
Name of antibody:	TNFSF12
Immunogen:	Synthesized peptide derived from internal of huma



www.shjning.com

	n TNF12.
Full name:	tumor necrosis factor (ligand) superfamily, member 12
Synonyms:	APO3L; DR3LG; TWEAK
SwissProt:	O43508
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	27 kDa
WB Positive control:	COLO cells, Huvec cells and 293 cells lysates
WB Recommended dilution:	500-3000
IF positive control:	Hela cells
IF Recommend dilution:	100-500









