

兔抗 TGFBI 多克隆抗体

中文名称: 兔抗 TGFBI 多克隆抗体

英文名称: Anti-TGFBI rabbit polyclonal antibody

别名: transforming growth factor beta induced; CSD; CDB1; CDG2; CSD1; CSD2; CSD3; EBMD; LCD1; BIGH3; CDGG1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: TGFBI

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes an RGD-containing protein that bin ds to type I, II and IV collagens. The RGD motif is fou nd in many extracellular matrix proteins modulating cel I adhesion and serves as a ligand recognition sequenc e for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochon drial bone formation in cartilage. The protein is induce d by transforming growth factor-beta and acts to inhib it cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy.
Applications:	ELISA, WB, IHC
Name of antibody:	TGFBI
Immunogen:	Fusion protein of human TGFBI



www.shjning.com

Full name:	transforming growth factor beta induced
Synonyms:	CSD; CDB1; CDG2; CSD1; CSD2; CSD3; EBMD; LCD1; BI GH3; CDGG1
SwissProt:	Q15582
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	75 kDa
WB Positive control:	Rat liver tissue lysate
WB Recommended dilution:	500-2000





