

兔抗 PLCG1 (Phospho-Tyr783)多克隆抗体

- 中文名称: 兔抗 PLCG1 (Phospho-Tyr783)多克隆抗体
- 英文名称: Anti-PLCG1 (Phospho-Tyr783) rabbit polyclonal antibody
- 别名: PLC1; NCKAP3; PLC-II; PLC148; PLCgamma1
- 相关类别: 一抗
- 储存: 冷冻 (-20℃) 避光
- 宿主: Rabbit
- 抗原: PLCG1 (Phospho-Tyr783)
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor Ras GRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.
Applications:	WB

Name of antibody:	PLCG1 (Phospho-Tyr783)
Immunogen:	Synthetic peptide of human PLCG1 (Phospho-Tyr783)
Full name:	phospholipase C, gamma 1 (Phospho-Tyr783)
Synonyms :	PLC1; NCKAP3; PLC-II; PLC148; PLCgamma1
SwissProt:	P19174
WB Predicted band size:	155 kDa
WB Positive control:	HL60 cells untreated or treated with EGF
WB Recommended dilution:	500-1000

