

兔抗 MAPK14 多克隆抗体

中文名称: 兔抗 MAPK14 多克隆抗体

英文名称: Anti-MAPK14 rabbit polyclonal antibody

别 名: mitogen-activated protein kinase 14; RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: MAPK14

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

AP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wi de variety of cellular processes such as proliferation, diffe rentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its pho sphorylation by MAP kinase kinases (MKKs), or its autoph osphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX,

The protein encoded by this gene is a member of the M

cell cycle regulator CDC25B, and tumor suppressor p53,



	which suggest the roles of this kinase in stress related tr anscription and cell cycle regulation, as well as in genoto xic stress response. Four alternatively spliced transcript va riants of this gene encoding distinct isoforms have been reported.
Applications:	ELISA, IHC
Name of antibody:	MAPK14
Immunogen:	Fusion protein of human MAPK14
Full name:	mitogen-activated protein kinase 14
Synonyms:	RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA
SwissProt:	Q16539
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	100-300

