

兔抗 SETD7 多克隆抗体

中文名称: 兔抗 SETD7 多克隆抗体

英文名称: Anti-SETD7 rabbit polyclonal antibody

别 名: KMT7; SET7; SET9; SET7/9

相关类别: 一抗

储 存: 冷冻(-20℃)

抗原: SETD7

宿 主: Rabbit

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

ression. Histone methylation can occur on Arg or L ys residues, with an exquisite site selectivity for Lys methylation at specific positions in the N-termini of histones H3 and H4. SET7/9, a histone methyltransf erase (HMTase), which transfers methyl groups to L ys4 of histone H3, forms a complex with S-adenosy I-L-methionine. This complex contains an active site consisting of a binding pocket where an AdoMet molecule in an unusual conformation binds, a narro w substrate-specific channel that only unmethylated

The methylation of histones plays a pivotal role in the regulation of chromatin structure and gene exp

lysine residues can access and a catalytic tyrosine r



	esidue.
Applications:	ELISA, IHC
Name of antibody:	SETD7
Immunogen:	Fusion protein of human SETD7
Full name:	SET domain containing (lysine methyltransferase) 7
Synonyms:	KMT7; SET7; SET9; SET7/9
SwissProt:	Q8WTS6
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human tonsil and human liver cancer
IHC Recommend dilution:	50-200

