

兔抗 NAT8 多克隆抗体

中文名称: 兔抗 NAT8 多克隆抗体

英文名称: Anti-NAT8 rabbit polyclonal antibody

别 名: GLA; CML1; Hcml1; ATase2; TSC501; TSC510

相关类别: 一抗

抗 原: NAT8

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene, isolated using the differential display method to detect tissue-specific genes, is specifically expressed in kidney and liver. The encoded protein shows amino a cid sequence similarity to N-acetyltransferases. A similar protein in Xenopus affects cell adhesion and gastrulation movements, and may be localized in the secretory pathway. A highly similar paralog is found in a cluster with this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	NAT8
Immunogen:	Fusion protein of human NAT8
Full name:	N-acetyltransferase 8 (GCN5-related, putative)
Synonyms:	GLA; CML1; Hcml1; ATase2; TSC501; TSC510
SwissProt:	Q9UHE5



www.shjning.com

ELISA Recommended dilution:	1000-5000
IHC positive control:	Human lung cancer and Human cervical cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	26 kDa
WB Positive control:	Mouse kidney and liver tissue
WB Recommended dilution:	200-1000







