

## 兔抗 TGM3 多克隆抗体

中文名称: 兔抗 TGM3 多克隆抗体

英文名称: Anti-TGM3 rabbit polyclonal antibody

别 名: transglutaminase 3; TGE; UHS2

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: TGM3

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

| Background:       | Transglutaminases are enzymes that catalyze the crosslin king of proteins by epsilon-gamma glutamyl lysine isope ptide bonds. While the primary structure of transglutami nases is not conserved, they all have the same amino a cid sequence at their active sites and their activity is cal cium-dependent. The protein encoded by this gene cons ists of two polypeptide chains activated from a single pr ecursor protein by proteolysis. The encoded protein is in volved the later stages of cell envelope formation in the epidermis and hair follicle. |
|-------------------|--|
| Applications:     | ELISA, WB, IHC   |
| Name of antibody: | TGM3   |
| Immunogen:        | Fusion protein of human TGM3   |
| Full name:        | transglutaminase 3   |
| Synonyms:         | TGE; UHS2  |



## www.shjning.com

| SwissProt:                  | Q08188                     |
|-----------------------------|----------------------------|
| ELISA Recommended dilution: | 5000-10000                 |
| IHC positive control:       | Human liver cancer         |
| IHC Recommend dilution:     | 50- 200                    |
| WB Predicted band size:     | 77 kDa                     |
| WB Positive control:        | Human tongue tissue lysate |
| WB Recommended dilution:    | 1000-5000                  |





