

兔抗 SLC13A3 多克隆抗体

中文名称: 兔抗 SLC13A3 多克隆抗体

英文名称: Anti-SLC13A3 rabbit polyclonal antibody

别 名: NADC3; SDCT2

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: SLC13A3

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Mammalian sodium-dicarboxylate cotransporters transp
ort succinate and other Krebs cycle intermediates. The
y fall into 2 categories based on their substrate affinit
y: low affinity and high affinity. Both the low- and hig
h-affinity transporters play an important role in the ha
ndling of citrate by the kidneys. The protein encoded
by this gene represents the high-affinity form. Alternat
ively spliced transcript variants encoding different isofo
rms have been found for this gene, although the full-l
ength nature of some of them have not been charact
erized yet.
ELISA, WB
SLC13A3



www.shjning.com

Immunogen:	Synthetic peptide of human SLC13A3
Full name:	solute carrier family 13 (sodium-dependent dicarboxyla te transporter), member 3
Synonyms:	NADC3; SDCT2
SwissProt:	Q8WWT9
ELISA Recommended dilution:	1000-2000
WB Predicted band size:	67 kDa
WB Positive control:	Human plasma and placenta tissue
WB Recommended dilution:	200-1000

