

## 兔抗 NCOR2 多克隆抗体

中文名称：兔抗 NCOR2 多克隆抗体

英文名称：Anti-NCOR2 rabbit polyclonal antibody

别名：SMRT; TRAC; CTG26; SMRTE; TRAC1; N-CoR2; TNRC14; TRAC-1; SMAP270; SMRTE-tau

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：NCOR2

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

|                      |  |
|----------------------|--|
| <b>Background:</b>   | This gene encodes a nuclear receptor co-repressor that mediates transcriptional silencing of certain target genes . The encoded protein is a member of a family of thyroid hormone- and retinoic acid receptor-associated co-repressors. This protein acts as part of a multisubunit complex which includes histone deacetylases to modify chromatin structure that prevents basal transcriptional activity of target genes. Aberrant expression of this gene is associated with certain cancers. Alternate splicing results in multiple transcript variants encoding different isoforms . |
| <b>Applications:</b> | ELISA, IHC   |

|                                    |  |
|------------------------------------|--|
| <b>Name of antibody:</b>           | NCOR2  |
| <b>Immunogen:</b>                  | Synthetic peptide of human NCOR2   |
| <b>Full name:</b>                  | nuclear receptor corepressor 2   |
| <b>Synonyms :</b>                  | SMRT; TRAC; CTG26; SMRTE; TRAC1; N-CoR2; TNRC14; T RAC-1; SMAP270; SMRTE-tau |
| <b>SwissProt:</b>                  | Q9Y618   |
| <b>ELISA Recommended dilution:</b> | 2000-5000  |
| <b>IHC positive control:</b>       | Human cervical cancer  |
| <b>IHC Recommend dilution:</b>     | 25-100   |

