

兔抗 ME3 多克隆抗体

中文名称: 兔抗 ME3 多克隆抗体

英文名称: Anti-ME3 rabbit polyclonal antibody

别 名: NADP-ME

储 存: 冷冻(-20℃) 避光

抗 原: ME3

宿 主: Rabbit

相关类别: 一抗

反应种属: Human Mouse

标 记 物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	Malic enzyme catalyzes the oxidative decarboxylation of malate to pyruvate using either NAD+ or NADP+ as a co factor. Mammalian tissues contain 3 distinct isoforms of malic enzyme: a cytosolic NADP(+)-dependent isoform, a mitochondrial NADP(+)-dependent isoform, and a mitochondrial NAD(+)-dependent isoform. This gene encodes a m itochondrial NADP(+)-dependent isoform. Multiple alternat ively spliced transcript variants have been found for this g ene, but the biological validity of some variants has not b een determined.
Applications:	WB
Name of antibody:	ME3



www.shjning.com

Immunogen:	Synthesized peptide derived from internal of human ME3.
Full name:	malic enzyme 3, NADP(+)-dependent, mitochondrial
Synonyms:	NADP-ME
SwissProt:	Q16798
WB Predicted band size:	67 kDa
WB Positive control:	NIH/3T3 cells and RAW264.7 cells lysates
WB Recommended dilution:	500-3000

