

兔抗 OSBPL2 多克隆抗体

中文名称：兔抗 OSBPL2 多克隆抗体

英文名称：Anti-OSBPL2 rabbit polyclonal antibody

别名：oxysterol binding protein like 2; ORP2; ORP-2; DFNA67; DNFA67

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：OSBPL2

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain, although the encoded protein contains only the sterol-binding domain. In vitro studies have shown that the encoded protein can bind strongly to phosphatidic acid and weakly to phosphatidylinositol 3-phosphate, but cannot bind to 25-hydroxycholesterol. The protein associates with the Golgi apparatus. Transcript variants encoding different isoforms have been described.
Applications:	ELISA, IHC
Name of antibody:	OSBPL2

Immunogen:	Fusion protein of human OSBPL2
Full name:	oxysterol binding protein like 2
Synonyms:	ORP2; ORP-2; DFNA67; DNFA67
SwissProt:	Q9H1P3
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil and Human cervical cancer
IHC Recommend dilution:	50-300



