

兔抗 PSMA3 多克隆抗体

- 中文名称：兔抗 PSMA3 多克隆抗体
- 英文名称：Anti-PSMA3 rabbit polyclonal antibody
- 别名：proteasome subunit alpha 3; HC8; PSC3
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：PSMA3
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Two alternative transcripts encoding

	g different isoforms have been identified.
Applications:	ELISA, WB, IHC
Name of antibody:	PSMA3
Immunogen:	Fusion protein of human PSMA3
Full name:	proteasome subunit alpha 3
Synonyms:	HC8; PSC3
SwissProt:	P25788
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human ovarian cancer and Human thyroid cancer
IHC Recommend dilution:	40-200
WB Predicted band size:	28 kDa
WB Positive control:	NIH/3T3, A549, HL60 and PC3 cell, Mouse liver tissue, Mouse spleen tissue, Hela cell lysates
WB Recommended dilution:	500-2000



