

## 兔抗 PCBP4 多克隆抗体

中文名称：兔抗 PCBP4 多克隆抗体

英文名称：Anti-PCBP4 rabbit polyclonal antibody

别名：poly(rC) binding protein 4; CBP; LIP4; MCG10

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：PCBP4

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>WB Recommended dilution:</b>	200-1000
<b>WB Positive control:</b>	Mouse kidney tissue
<b>WB Predicted band size:</b>	41 kDa
<b>SwissProt:</b>	P57723
<b>Synonyms:</b>	CBP; LIP4; MCG10
<b>Full name:</b>	poly(rC) binding protein 4
<b>Immunogen:</b>	Synthetic peptide of human PCBP4
<b>Name of antibody:</b>	PCBP4
<b>Applications:</b>	ELISA, WB
<b>Background:</b>	This gene encodes a member of the KH-domain protein subfamily. Proteins of this subfamily, also referred to as alpha-CPs, bind to RNA with a specificity for C-rich pyrimidin

e regions. Alpha-CPs play important roles in post-transcriptional activities and have different cellular distributions. This gene is induced by the p53 tumor suppressor, and the encoded protein can suppress cell proliferation by inducing apoptosis and cell cycle arrest in G(2)-M. This gene's protein is found in the cytoplasm, yet it lacks the nuclear localization signals found in other subfamily members. Multiple alternatively spliced transcript variants have been described, but the full-length nature for only some has been determined. [provided by RefSeq, Jul 2008]

