

## 兔抗 NFATC2 多克隆抗体

中文名称: 兔抗 NFATC2 多克隆抗体

英文名称: Anti-NFATC2 rabbit polyclonal antibody

别 名: NFAT1, NFATP

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

克隆类型: rabbit polyclonal

抗 原: NFATC2

反应种属: Human, Mouse

标 记 物: Unconjugate

## 技术规格

Name of antibody:	NFATC2
Applications:	ELISA, WB
	have been characterized.
Background:	transcriptional splice variants encoding different isoforms
	gene transcription during the immune response. Alternate
	on complex. This complex plays a central role in inducing
	mber of the nuclear factors of activated T cells transcripti
	T cell receptor (TCR) stimulation, where it becomes a me
	n the cytosol and only translocates to the nucleus upon
	n NFAT-homology region (NHR). This protein is present i
	binding protein with a REL-homology region (RHR) and a
	T cells (NFAT) family. The product of this gene is a DNA-
	This gene is a member of the nuclear factor of activated



Immunogen:	Synthetic peptide of human NFATC2
Full name:	Nuclear factor of activated T-cells, cytoplasmic, calcineurin -dependent 2
Synonyms:	NFAT1, NFATP
SwissProt:	Q13469
ELISA Recommended dilution:	1000-2000
WB Predicted band size:	98 kDa
WB Positive control:	Human fetal brain tissue
WB Recommended dilution:	200-1000

