

兔抗 TNFRSF8 多克隆抗体

中文名称：兔抗 TNFRSF8 多克隆抗体

英文名称：Anti-TNFRSF8 rabbit polyclonal antibody

别名：CD30; Ki-1; D1S166E

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：TNFRSF8

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. |
| Applications: | ELISA, IHC |
| Name of antibody: | TNFRSF8 |
| Immunogen: | Synthetic peptide of human TNFRSF8 |

| | |
|------------------------------------|--|
| Full name: | tumor necrosis factor receptor superfamily, member 8 |
| Synonyms : | CD30; Ki-1; D1S166E |
| SwissProt: | P28908 |
| ELISA Recommended dilution: | 1000-5000 |
| IHC positive control: | Human Lymphoma and Human thyroid cancer |
| IHC Recommend dilution: | 50-200 |

