

## 兔抗 NECTIN4 多克隆抗体

中文名称: 兔抗 NECTIN4 多克隆抗体

英文名称: Anti-NECTIN4 rabbit polyclonal antibody

别 名: nectin cell adhesion molecule 4; LNIR; PRR4; EDSS1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: NECTIN4

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

ed protein contains two immunoglobulin-like (Ig-like) C2-type domains and one Ig-like V-type domain. It is involved in cell adhesion through trans-homophilic and -heterophilic interacti ons. It is a single-pass type I membrane protein. The soluble form is produced by proteolytic cleavage at the cell surface by the metalloproteinase ADAM17/TACE. The secreted form is found in both breast tumor cell lines and breast tumor patients. Mutations in this gene are the cause of ectodermal dyspl asia-syndactyly syndrome type 1, an autosomal recessive disor der. Alternatively spliced transcript variants have been found but the full-length nature of the variant has not been determ

ined.[provided by RefSeq, Jan 2011]

This gene encodes a member of the nectin family. The encod

## **Background:**



| Applications:           | ELISA, IHC                                |
|-------------------------|---|
| Name of antibody:       | NECTIN4                                   |
| Immunogen:              | Fusion protein of human NECTIN4           |
| Full name:              | nectin cell adhesion molecule 4           |
| Synonyms:               | LNIR; PRR4; EDSS1; PVRL4; nectin-4        |
| SwissProt:              | Q96NY8                                    |
| IHC positive control:   | Human colon cancer and Human liver cancer |
| IHC Recommend dilution: | 50-200                                    |

