

## 兔抗 GTF2I 多克隆抗体

中文名称：兔抗 GTF2I 多克隆抗体

英文名称：Anti-GTF2I rabbit polyclonal antibody

别名：WBS; DIWS; SPIN; IB291; BAP135; BTKAP1; TFII-I; WBSCR6; GTFII-I

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：GTF2I

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene encodes a phosphoprotein containing six characteristic repeat motifs. The encoded protein binds to the initiator element (Inr) and E-box element in promoters and functions as a regulator of transcription. This locus, along with several other neighboring genes, is deleted in Williams-Beuren syndrome. There are many closely related genes and pseudogenes for this gene on chromosome 7. This gene also has pseudogenes on chromosomes 9, 13, and 21. Alternatively spliced transcript variants encoding multiple isoforms have been observed.
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	GTF2I
<b>Immunogen:</b>	Synthetic peptide of human GTF2I
<b>Full name:</b>	general transcription factor Ii
<b>Synonyms :</b>	WBS; DIWS; SPIN; IB291; BAP135; BTKAP1; TFII-I; W BSCR6; GTFII-I
<b>SwissProt:</b>	P78347
<b>ELISA Recommended dilution:</b>	1000-2000
<b>IHC positive control:</b>	Human thyroid cancer and Human cervical cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	112 kDa
<b>WB Positive control:</b>	HepG2 cells
<b>WB Recommended dilution:</b>	200-1000



