

兔抗 PSMD2 多克隆抗体

中文名称：兔抗 PSMD2 多克隆抗体

英文名称：Anti-PSMD2 rabbit polyclonal antibody

别名：S2; P97; Rpn1; TRAP2

抗原：PSMD2

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, th
--------------------	--

	<p>e immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator lid. In addition to participation in proteasome function, this subunit may also participate in the TNF signalling pathway since it interacts with the tumor necrosis factor type 1 receptor. A pseudogene has been identified on chromosome 1.</p>
Applications:	ELISA, WB
Name of antibody:	PSMD2
Immunogen:	Fusion protein of human PSMD2
Full name:	Proteasome (prosome, macropain) 26S subunit, non-ATPase, 2
Synonyms :	S2; P97; Rpn1; TRAP2
SwissProt:	Q13200
ELISA Recommended dilution:	1000-2000
WB Predicted band size:	100 kDa
WB Positive control:	Jurkat cells
WB Recommended dilution:	200-1000

