

## 兔抗 SLC18A2 多克隆抗体

中文名称：兔抗 SLC18A2 多克隆抗体

英文名称：Anti-SLC18A2 rabbit polyclonal antibody

别名：SVAT, SVMT, VAT2, VMAT2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC18A2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	The vesicular monoamine transporter acts to accumulate cytosolic monoamines into synaptic vesicles, using the proton gradient maintained across the synaptic vesicular membrane. Its proper function is essential to the correct activity of the monoaminergic systems that have been implicated in several human neuropsychiatric disorders. The transporter is a site of action of important drugs, including reserpine and tetrabenazine .
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	SLC18A2
<b>Immunogen:</b>	Synthetic peptide of human SLC18A2
<b>Full name:</b>	solute carrier family 18 (vesicular monoamine), me

	mber 2
<b>Synonyms :</b>	SVAT, SVMT, VAT2, VMAT2
<b>SwissProt:</b>	Q05940
<b>ELISA Recommended dilution:</b>	2000-60000
<b>IHC positive control:</b>	Human brain
<b>IHC Recommend dilution:</b>	25-100
<b>WB Predicted band size:</b>	55 kDa
<b>WB Positive control:</b>	Human cervical cancer tissue
<b>WB Recommended dilution:</b>	500-2000



