

兔抗 SLC5A10 多克隆抗体

中文名称：兔抗 SLC5A10 多克隆抗体

英文名称：Anti-SLC5A10 rabbit polyclonal antibody

别名：SGLT5; SGLT-5

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC5A10

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene is a member of the sodium/glucose transporter family. Members of this family are sodium-dependent transporters and can be divided into two subfamilies based on sequence homology, one that co-transportes sugars and the second that transports molecules such as ascorbate, choline, iodide, lipolate, monocarboxylates, and pantothenate. The protein encoded by this gene has the highest affinity for mannose and has been reported to be most highly expressed in the kidney. This protein may function as a kidney-specific, sodium-dependent mannose and fructose co-transporter. Alternative splicing results in multiple transcript variants that encode different protein isoforms.
Applications:	ELISA, IHC

Name of antibody:	SLC5A10
Immunogen:	Synthetic peptide of human SLC5A10
Full name:	solute carrier family 5 (sodium/sugar cotransporter), member 10
Synonyms:	SGLT5; SGLT-5
SwissProt:	A0PJK1
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human brain and human colorectal cancer
IHC Recommend dilution:	20-100



