

## 兔抗 HADHA 多克隆抗体

- 中文名称：兔抗 HADHA 多克隆抗体
- 英文名称：Anti-HADHA rabbit polyclonal antibody
- 别名：GBP; ECHA; HADH; LCEH; MTPA; LCHAD; TP-ALPHA
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：HADHA
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 相关类别：一抗
- 克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene encodes the alpha subunit of the mitochondrial trifunctional protein, which catalyzes the last three steps of mitochondrial beta-oxidation of long chain fatty acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four beta subunits, with the alpha subunit catalyzing the 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratase activities. Mutations in this gene result in trifunctional protein deficiency or LCHAD deficiency. The genes of the alpha and beta subunits of the mitochondrial trifunctional protein are located adjacent to each other in the human genome in a head-to-head orientation.
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	HADHA
<b>Immunogen:</b>	Fusion protein of human HADHA
<b>Full name:</b>	hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase /enoyl-CoA hydratase (trifunctional protein), alpha subunit
<b>Synonyms:</b>	GBP; ECHA; HADH; LCEH; MTPA; LCHAD; TP-ALPHA
<b>SwissProt:</b>	P40939
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human ovarian cancer
<b>IHC Recommend dilution:</b>	25-100
<b>WB Predicted band size:</b>	83 kDa
<b>WB Positive control:</b>	A431 , HEPG2 , Jurkat and 293T cell lysates
<b>WB Recommended dilution:</b>	500-2000



