

兔抗 STARD13 多克隆抗体

中文名称：兔抗 STARD13 多克隆抗体

英文名称： Anti-STARD13 rabbit polyclonal antibody

别名： DLC2; GT650; ARHGAP37; LINC00464

抗原： STARD13

储存： 冷冻（-20℃） 避光

宿主： Rabbit

反应种属： Human

相关类别： 一抗

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

Background:	This gene encodes a protein which contains an N-terminal sterile alpha motif (SAM) for protein-protein interactions, followed by an ATP/GTP-binding motif, a GTPase-activating protein (GAP) domain, and a C-terminal STAR-related lipid transfer (START) domain. It may be involved in regulation of cytoskeletal reorganization, cell proliferation, and cell motility, and acts as a tumor suppressor in hepatoma cells. The gene is located in a region of chromosome 13 that is associated with loss of heterozygosity in hepatocellular carcinomas. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
Applications:	WB, IF
Name of antibody:	STARD13
Immunogen:	Synthesized peptide derived from N-terminal of human STARD13.

Full name:	StAR-related lipid transfer (START) domain containing 13
Synonyms :	DLC2; GT650; ARHGAP37; LINC00464
SwissProt:	Q9Y3M8
WB Predicted band size:	125 kDa
WB Positive control:	K562 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	A549 cells
IF Recommend dilution:	100-500



