

兔抗 TNK2(Phospho-Tyr284) 多克隆抗体

中文名称：兔抗 TNK2(Phospho-Tyr284) 多克隆抗体

英文名称：Anti-TNK2(Phospho-Tyr284) rabbit polyclonal antibody

别名：ACK; ACK1; ACK-1; p21cdc42Hs

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：TNK2(Phospho-Tyr284)

反应种属：Human Mouse

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined.
Applications:	WB, IHC, IF
Name of antibody:	TNK2(Phospho-Tyr284)
Immunogen:	Peptide sequence around phosphorylation site of tyrosin

	e 284 (D-H-Y(p)-V-M) derived from Human ACK1.
Full name:	tyrosine kinase, non-receptor, 2
Synonyms :	ACK; ACK1; ACK-1; p21cdc42Hs
SwissProt:	Q07912
IHC positive control:	Human breast carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	115 kDa
WB Positive control:	HepG2 cells lysates treated with EGF
WB Recommended dilution:	500-1000
IF positive control:	A549 cells
IF Recommend dilution:	100-200



