

兔抗 MET 多克隆抗体

| | |
|-------|-------------------------------------|
| 中文名称: | 兔抗 MET 多克隆抗体 |
| 英文名称: | Anti-MET rabbit polyclonal antibody |
| 别名: | HGFR; AUTS9; RCCP2; c-Met |
| 相关类别: | 一抗 |
| 储存: | 冷冻 (-20℃) |
| 宿主: | Rabbit |
| 抗原: | MET |
| 反应种属: | Human, Mouse, Rat |
| 标记物: | Unconjugate |
| 克隆类型: | rabbit polyclonal |

技术规格

| | |
|--------------------------|--|
| Background: | The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma. Two transcript variants encoding different isoforms have been found for this gene. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | MET |
| Immunogen: | Synthetic peptide of human MET |
| Full name: | MET proto-oncogene, receptor tyrosine kinase |
| Synonyms: | HGFR; AUTS9; RCCP2; c-Met; DFN97 |

| | |
|--------------------------|----------------------------|
| SwissProt: | P08581 |
| IHC positive control: | Human esophagus cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 156 kDa |
| WB Positive control: | Hela and A431 cell lysates |
| WB Recommended dilution: | 500-2000 |



