

兔抗 RPL30 多克隆抗体

中文名称: 兔抗 RPL30 多克隆抗体

英文名称: Anti-RPL30 rabbit polyclonal antibody

别 名: L30

抗 原: RPL30

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

反应种属: Human Mouse

相关类别: 一抗

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	Ribosomes, the organelles that catalyze protein syn thesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are compose d of 4 RNA species and approximately 80 structura lly distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L30E family of ribosomal proteins. It is located in the cytoplasm. This gene is co-transcribed with the U72 small nucleolar RNA gene, which is located in its fourth intron. As is ty pical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene d ispersed through the genome.
Applications:	WB, IHC



www.shjning.com

Name of antibody:	RPL30
Immunogen:	Synthesized peptide derived from internal of huma n RPL30.
Full name:	ribosomal protein L30
Synonyms:	L30
SwissProt:	P62888
IHC positive control:	Human prostate carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	13 kDa
WB Positive control:	A549 cells, COS-7 cells and RAW264.7 cells lysates
WB Recommended dilution:	500-3000





