

兔抗 TAF11 多克隆抗体

中文名称：兔抗 TAF11 多克隆抗体

英文名称： Anti-TAF11 rabbit polyclonal antibody

别名： TAF2I; PRO2134; TAFII28; MGC:15243

相关类别： 一抗

储存： 冷冻 (-20℃)

宿主： Rabbit

抗原： TAF11

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit of TFIID that is present in all TFIID complexes and interacts with TBP. This subunit also interacts with another

	her small subunit, TAF13, to form a heterodimer with a structure similar to the histone core structure. Two transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	TAF11
Immunogen:	Fusion protein of human TAF11
Full name:	TAF11 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 28kDa
Synonyms :	TAF2I; PRO2134; TAFII28; MGC:15243
SwissProt:	Q15544
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	35-150
WB Predicted band size:	23 kDa
WB Positive control:	Mouse brain and heart tissue, Jurkat, K562 and hela cells
WB Recommended dilution:	500-2000



