

兔抗 TAF6L 多克隆抗体

中文名称：兔抗 TAF6L 多克隆抗体

英文名称：Anti-TAF6L rabbit polyclonal antibody

别名：PAF65A

抗原：TAF6L

储存：冷冻（-20℃）避光

宿主：Rabbit

反应种属：Human Mouse

相关类别：一抗

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a protein that is a component of the PCAF histone acetylase complex and structurally similar to one of the histone-like TAFs, TAF6. The PCAF
--------------------	--

	histone acetylase complex, which is composed of more than 20 polypeptides some of which are TAFs, is required for myogenic transcription and differentiation.
Applications:	WB
Name of antibody:	TAF6L
Immunogen:	Synthesized peptide derived from internal of human TAF6L.
Full name:	TAF6-like RNA polymerase II, p300/CBP-associated factor (P CAF)-associated factor, 65kDa
Synonyms :	PAF65A
SwissProt:	Q9Y6J9
WB Predicted band size:	68 kDa
WB Positive control:	293 cells lysate
WB Recommended dilution:	500-3000

