

兔抗 RPL36 多克隆抗体

- 中文名称: 兔抗 RPL36 多克隆抗体
- 英文名称: Anti-RPL36 rabbit polyclonal antibody
- 别名: L36
- 抗原: RPL36
- 储存: 冷冻(-20℃) 避光
- 宿 主: Rabbit
- 反应种属: Human
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	Ribosomes, the organelles that catalyze protein synth esis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distince t proteins. This gene encodes a ribosomal protein th at is a component of the 60S subunit. The protein b elongs to the L36E family of ribosomal proteins. It is located in the cytoplasm. Transcript variants derived the rom alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal pro- teins, there are multiple processed pseudogenes of this gene dispersed through the genome.
Applications:	WB, IHC, IF
	this gene dispersed through the genome.
Background:	roteins, there are multiple processed pseudogenes of
	protein. As is typical for genes encoding ribosomal p
	rom alternative splicing exist; they encode the same
	located in the cytoplasm. Transcript variants derived
	esis, consist of a small 40S subunit and a large 60S
	Ribosomes, the organelles that catalyze protein synth



Name of antibody:	RPL36
Immunogen:	Synthesized peptide derived from internal of human RPL36.
Full name:	ribosomal protein L36
Synonyms :	L36
SwissProt:	Q9Y3U8
IHC positive control:	Human prostate carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	12 kDa
WB Positive control:	COLO cells lysate
WB Recommended dilution:	500-3000
IF positive control:	A549 cells
IF Recommend dilution:	100-500







