

www.shjning.com

兔抗 RIDA 多克隆抗体

- 中文名称: 兔抗 RIDA 多克隆抗体
- 英文名称: Anti-RIDA rabbit polyclonal antibody

别名: reactive intermediate imine deaminase A homolog; PSP; P14.5; UK114; HRSP12; hp14.5

- 相关类别: 一抗
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗原: RIDA
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	Catalyzes the hydrolytic deamination of enamine/imin
	e intermediates that form during the course of norm
	al metabolism. May facilitate the release of ammonia
	from these potentially toxic reactive metabolites, redu
	cing their impact on cellular components. It may act
	on enamine/imine intermediates formed by several ty
	pes of pyridoxal-5'-phosphate-dependent dehydratase
	s including L-threonine dehydratase. May also functio
	n as an endoribonuclease, cleaving mRNA phosphodi
	ester bonds of single-stranded RNA (By similarity). Th
	ereby, may inhibit protein translation (PubMed:897365
	3).



Applications:	ELISA, WB, IHC
Name of antibody:	RIDA
Immunogen:	Fusion protein of human RIDA
Full name:	reactive intermediate imine deaminase A homolog
Synonyms:	PSP; P14.5; UK114; HRSP12; hp14.5
SwissProt:	P52758
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human lung cancer and Human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	14 kDa
WB Positive control:	Rat kidney tissue, Mouse liver tissue and Human fetal
	liver tissue lysates
WB Recommended dilution:	500-2000





