

兔抗 PLD2(Phospho-Tyr169) 多克隆抗体

中文名称：兔抗 PLD2(Phospho-Tyr169) 多克隆抗体

英文名称：Anti-PLD2(Phospho-Tyr169) rabbit polyclonal antibody

别名：PLD2

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：PLD2(Phospho-Tyr169)

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | The protein encoded by this gene catalyzes the hydrolysis of phosphatidylcholine to phosphatidic acid and choline. The activity of the encoded enzyme is enhanced by phosphatidylinositol 4,5-bisphosphate and ADP-ribosylation factor-1. This protein localizes to the peripheral membrane and may be involved in cytoskeletal organization, cell cycle control, transcriptional regulation, and/or regulated secretion. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011] |
| Applications: | WB, IHC |
| Name of antibody: | PLD2(Phospho-Tyr169) |
| Immunogen: | Peptide sequence around phosphorylation site of tyrosine 1 |

| | |
|---------------------------------|---|
| | 69(E-N-Y(p)-L-N) derived from Human PLD2. |
| Full name: | phospholipase D2 |
| Synonyms : | PLD2 |
| SwissProt: | O14939 |
| IHC positive control: | Human brain tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 106 kDa |
| WB Positive control: | Jurkat cells lysates treated with TNF |
| WB Recommended dilution: | 500-1000 |



