

兔抗 STRADB 多克隆抗体

中文名称: 兔抗 STRADB 多克隆抗体

英文名称: Anti-STRADB rabbit polyclonal antibody

别 名: PAPK; ILPIP; ILPIPA; ALS2CR2; CALS-21; PRO1038

相关类别: 一抗

储 存: 冷冻(-20℃)

抗 原: STRADB

宿 主: Rabbit

反应种属: Human, Mouse

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

This gene encodes a protein that belongs to the serin e/threonine protein kinase STE20 subfamily. One of the active site residues in the protein kinase domain of this protein is altered, and it is thus a pseudokinase. This protein is a component of a complex involved in the activation of serine/threonine kinase 11, a master kinase that regulates cell polarity and energy-generating metabolism. This complex regulates the relocation of this kinase from the nucleus to the cytoplasm, and it is essential for G1 cell cycle arrest mediated by this kinase. The protein encoded by this gene can also interact with the X chromosome-linked inhibitor of apop tosis protein, and this interaction enhances the anti-a poptotic activity of this protein via the JNK1 signal transduction pathway. Two pseudogenes, located on chr



www.shjning.com

	omosomes 1 and 7, have been found for this gene. A Iternatively spliced transcript variants encoding differe nt isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	STRADB
Immunogen:	Fusion protein of human STRADB
Full name:	STE20-related kinase adaptor beta
Synonyms:	PAPK; ILPIP; ILPIPA; ALS2CR2; CALS-21; PRO1038
SwissProt:	Q9C0K7
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human lymphoma and Human thyroid cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	47 kDa; 31 kDa
WB Positive control:	Mouse pancreas and spleen tissue, mouse heart tissue and 293T cells
WB Recommended dilution:	200-1000





