

兔抗 SLC22A12 多克隆抗体

中文名称: 兔抗 SLC22A12 多克隆抗体

英文名称: Anti-SLC22A12 rabbit polyclonal antibody

别 名: solute carrier family 22 member 12; RST; OAT4L; URAT1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: SLC22A12

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the o rganic anion transporter (OAT) family, and it acts as a ur ate transporter to regulate urate levels in blood. This pr otein is an integral membrane protein primarily found in epithelial cells of the proximal tubule of the kidney. An elevated level of serum urate, hyperuricemia, is associate d with increased incidences of gout, and mutations in th is gene cause renal hypouricemia type 1. Alternative spli cing results in multiple transcript variants.
Applications:	ELISA, WB, IHC
Name of antibody:	SLC22A12
Immunogen:	Synthetic peptide of human SLC22A12
Full name:	solute carrier family 22 member 12



www.shjning.com

Synonyms:	RST; OAT4L; URAT1
SwissProt:	Q96S37
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human cervical cancer and Human colorectal cancer
IHC Recommend dilution:	50-100
WB Predicted band size:	60 kDa
WB Positive control:	293T cell lysate
WB Recommended dilution:	500-2000







