

兔抗 MTIF2 多克隆抗体

中文名称：兔抗 MTIF2 多克隆抗体

英文名称：Anti-MTIF2 rabbit polyclonal antibody

别 名：mitochondrial translational initiation factor 2

相关类别：一抗

储 存：冷冻 (-20°C)

宿 主：Rabbit

抗 原：MTIF2

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格


Background:	During the initiation of protein biosynthesis, initiation factor-2 (IF-2) promotes the binding of the initiator tRNA to the small subunit of the ribosome in a GTP-dependent manner. Prokaryotic IF-2 is a single polypeptide, while eukaryotic cytoplasmic IF-2 (eIF-2) is a trimeric protein. Bovine liver mitochondria contain IF-2(mt), an 85-kD monomeric protein that is equivalent to prokaryotic IF-2. The predicted 727-amino acid human protein contains a 29-amino acid presequence. Human IF-2(mt) shares 32 to 38 % amino acid sequence identity with yeast IF-2(mt) and several prokaryotic IF-2s, with the greatest degree of conservation in the G domains of the proteins.
Applications:	ELISA, IHC



上海纪宁

www.shjning.com

Name of antibody:	MTIF2
Immunogen:	Synthetic peptide of human MTIF2
Full name:	mitochondrial translational initiation factor 2
SwissProt:	P46199
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil and Human breast cancer
IHC Recommend dilution:	20-100





上海纪宁

www.shjning.com

