

兔抗 TBCA 多克隆抗体

中文名称：兔抗 TBCA 多克隆抗体

英文名称： Anti-TBCA rabbit polyclonal antibody

相关类别： 一抗

储 存： 冷冻（-20℃）

宿 主： Rabbit

抗 原： TBCA

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	The product of this gene is one of four proteins (cofactors A, D, E, and C) involved in the pathway leading to correctly folded beta-tubulin from folding intermediates. Cofactors A and D are believed to play a role in capturing and stabilizing beta-tubulin intermediates in a quasi-native confirmation. Cofactor E binds to the cofactor D /beta-tubulin complex; interaction with cofactor C then causes the release of beta-tubulin polypeptides that are committed to the native state.
Applications:	ELISA, WB, IHC
Name of antibody:	TBCA
Immunogen:	Full length fusion protein
Full name:	tubulin folding cofactor A
SwissProt:	O75347
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human gastric cancer and human thyroid cancer

IHC Recommend dilution:	25-100
WB Predicted band size:	13 kDa
WB Positive control:	Mouse adrenal gland tissue
WB Recommended dilution:	200-1000



