

兔抗 SOCS2 多克隆抗体

中文名称：兔抗 SOCS2 多克隆抗体

英文名称：Anti-SOCS2 rabbit polyclonal antibody

别名：CIS2; SSI2; Cish2; SSI-2; SOCS-2; STAT12

抗原：SOCS2

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the suppressor of cytokine signaling (SOCS) family. SOCS family members are cytokine-inducible negative regulators of cytokine receptor signaling via the Janus kinase/signal transducer and activation of transcription pathway (the JAK/STAT pathway). SOCS family proteins interact with major molecules of signaling complexes to block further signal transduction, in part, by proteasomal depletion of receptors or signal-transducing proteins via ubiquitination. The expression of this gene can be induced by a subset of cytokines, including erythropoietin, GM-CSF, IL10, interferon (IFN)-gamma and by cytokine receptors such as growth hormone receptor.
Applications:	ELISA, IHC
Name of antibody:	SOCS2

Immunogen:	Fusion protein of human SOCS2
Full name:	suppressor of cytokine signaling 2
Synonyms :	CIS2; SSI2; Cish2; SSI-2; SOCS-2; STATI2
SwissProt:	O14508
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human brain
IHC Recommend dilution:	25-100

