

兔抗 SMOC2 多克隆抗体

中文名称: 兔抗 SMOC2 多克隆抗体

英文名称: Anti-SMOC2 rabbit polyclonal antibody

别 名: DTDP1; SMAP2; MST117; MSTP117; MSTP140; bA37D8.1; bA270C4A.1; dJ421D16.1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: SMOC2

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

| Background: | This gene encodes a member of the SPARC family (s ecreted protein acidic and rich in cysteine/osteonecti n/BM-40), which are highly expressed during embryo genesis and wound healing. The gene product is a matricellular protein which promotes matrix assembly and can stimulate endothelial cell proliferation and migration, as well as angiogenic activity. Associated with pulmonary function, this secretory gene product contains a Kazal domain, two thymoglobulin type-1 domains, and two EF-hand calcium-binding domains. |
|-------------------|---|
| Applications: | ELISA, WB, IHC |
| Name of antibody: | SMOC2 |
| Immunogen: | Synthetic peptide of human SMOC2 |
| Full name: | SPARC related modular calcium binding 2 |
| Synonyms : | DTDP1; SMAP2; MST117; MSTP117; MSTP140; bA37D |



www.shjning.com

| | 8.1; bA270C4A.1; dJ421D16.1 |
|-----------------------------|--|
| SwissProt: | Q9H3U7 |
| ELISA Recommended dilution: | 1000-2000 |
| IHC positive control: | Human cervical cancer and Human liver cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 50 kDa |
| WB Positive control: | Human placenta tissue |
| WB Recommended dilution: | 200-1000 |





