

兔抗 MED1(Phospho-Thr1457) 多克隆抗体

中文名称: 兔抗 MED1(Phospho-Thr1457) 多克隆抗体

英文名称: Anti-MED1(Phospho-Thr1457) rabbit polyclonal antibody

别名: PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220

相关类别: 一抗

储存: 冷冻 (-20°C) 避光

宿主: Rabbit

抗原: MED1(Phospho-Thr1457)

反应种属: Human Mouse

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

| | |
|--------------------------|---|
| Background: | The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. |
| Applications: | WB |
| Name of antibody: | MED1(Phospho-Thr1457) |
| Immunogen: | Peptide sequence around phosphorylation site of Threonine 1457 (A-Y-T(p)-P-Q) derived from Human PPAR-BP. |
| Full name: | mediator complex subunit 1 |

| | |
|---------------------------------|---|
| Synonyms : | PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220 |
| SwissProt: | Q15648 |
| WB Predicted band size: | 168 kDa |
| WB Positive control: | Huvec cells lysates treated with serum |
| WB Recommended dilution: | 500-1000 |

