

## 兔抗 GSTM4 多克隆抗体

- 中文名称: 兔抗 GSTM4 多克隆抗体
- 英文名称: Anti-GSTM4 rabbit polyclonal antibody
- 别 名: GTM4; GSTM4-4
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 抗 原: GSTM4
- 宿 主: Rabbit
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

## 技术规格

Background:	Cytosolic and membrane-bound forms of glutathion e S-transferase are encoded by two distinct superg ene families. At present, eight distinct classes of th e soluble cytoplasmic mammalian glutathione S-tran sferases have been identified: alpha, kappa, mu, om ega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu cl ass. The mu class of enzymes functions in the deto xification of electrophilic compounds, including carci nogens, therapeutic drugs, environmental toxins an d products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of e
-------------	---



	nzymes are organized in a gene cluster on chromo some 1p13.3 and are known to be highly polymorp hic. These genetic variations can change an individu al's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Diversification of these genes has occurred in regio ns encoding substrate-binding domains, as well as i n tissue expression patterns, to accommodate an in creasing number of foreign compounds. Multiple tr anscript variants, each encoding a distinct protein is oform, have been identified.
Applications:	elisa, wb, ihc
Name of antibody:	GSTM4
Immunogen:	Full length fusion protein
Full name:	glutathione S-transferase mu 4
Synonyms :	GTM4; GSTM4-4
SwissProt:	Q03013
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer and human lung cancer
IHC Recommend dilution:	10-50
WB Predicted band size:	26 kDa
WB Positive control:	Mouse liver tissue , Mouse lung tissue
WB Recommended dilution:	200-1000













