

兔抗 RBM8A 多克隆抗体

中文名称：兔抗 RBM8A 多克隆抗体

英文名称：Anti-RBM8A rabbit polyclonal antibody

别名：TAR; Y14; RBM8; ZNRP; RBM8B; ZRNP1; BOV-1A; BOV-1B; BOV-1C; MDS014; DEL1q21.1; C1DELq21.1

抗原：RBM8A

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a protein with a conserved RNA-binding motif. The protein is found predominantly in the nucleus, although it is also present in the cytoplasm. It is preferentially associated with mRNAs produced by splicing, including both nuclear mRNAs and newly exported cytoplasmic mRNAs. It is thought that the protein remains associated with spliced mRNAs as a tag to indicate where introns had been present, thus coupling pre- and post-mRNA splicing events. Previously, it was thought that two genes encode this protein, RBM8A and RBM8B; it is now thought that

	<p>t the RBM8B locus is a pseudogene. There are two alternate translation start codons with this gene, which result in two forms of the protein. An allele mutation and a low-frequency noncoding single-nucleotide polymorphism (SNP) in this gene cause thrombocytopenia-absent radius (TAR) syndrome.</p>
Applications:	ELISA, IHC
Name of antibody:	RBM8A
Immunogen:	Full length fusion protein
Full name:	RNA binding motif protein 8A
Synonyms:	TAR; Y14; RBM8; ZNRP; RBM8B; ZRNP1; BOV-1A; BOV-1B; BOV-1C; MDS014; DEL1q21.1; C1DELq21.1
SwissProt:	Q9Y5S9
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer
IHC Recommend dilution:	25-100



