

兔抗 TNFRSF17 多克隆抗体

- 中文名称: 兔抗 TNFRSF17 多克隆抗体
- 英文名称: Anti-TNFRSF17 rabbit polyclonal antibody
- 名: BCM; BCMA; CD269; TNFRSF13A 别
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: TNFRSF17
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be import ant for B cell development and autoimmune response. This receptor has been shown to specifically bind to th e tumor necrosis factor (ligand) superfamily, member 13 b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB a nd MAPK8/JNK activation. This receptor also binds to v arious TRAF family members, and thus may transduce si gnals for cell survival and proliferation
Applications:	ELISA, WB, IHC
Name of antibody:	TNFRSF17
Immunogen:	Synthetic peptide of human TNFRSF17
Full name:	tumor necrosis factor receptor superfamily, member 17
Synonyms:	BCM; BCMA; CD269; TNFRSF13A



www.shjning.com

SwissProt:	Q02223
IHC positive control:	Human gastric cancer and human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	20 kDa
WB Positive control:	Mouse skeletal muscle tissue
WB Recommended dilution:	200-1000





