

兔抗 NFATC1 多克隆抗体

- 中文名称：兔抗 NFATC1 多克隆抗体
- 英文名称：Anti-NFATC1 rabbit polyclonal antibody
- 别名：NFAT2, NFATc, NF-ATC
- 抗原：NFATC1
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 相关类别：一抗
- 反应种属：Human, Mouse
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

The product of this gene is a component of the nuclear factor of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation, and an inducible nuclear component. Proteins belonging to this family of transcription factors play a central role in inducible gene transcription during immune response. The product of this gene is an inducible nuclear component. It functions as a major molecular target for the immunosuppressive drugs such as cyclosporin A. Five transcript variants encoding distinct isoforms have been identified for this gene. Different isoforms of this protein may regulate inducible expression of different cytokine genes.

Applications:	ELISA, IHC
Name of antibody:	NFATC1
Immunogen:	Fusion protein of human NFATC1
Full name:	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1
Synonyms :	NFAT2, NFATc, NF-ATC
SwissProt:	O95644
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human esophagus cancer and Human cervical cancer
IHC Recommend dilution:	25-100

