

兔抗 TBCC 多克隆抗体

中文名称: 兔抗 TBCC 多克隆抗体

英文名称: Anti-TBCC rabbit polyclonal antibody

别 名: tubulin folding cofactor C; CFC

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: TBCC

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

| Background: | Cofactor C is one of four proteins (cofactors A, D, E, an d C) involved in the pathway leading to correctly folded beta-tubulin from folding intermediates. Cofactors A and D are believed to play a role in capturing and stabilizing beta-tubulin intermediates in a quasi-native confirmation. Cofactor E binds to the cofactor D/beta-tubulin complex; interaction with cofactor C then causes the release of be ta-tubulin polypeptides that are committed to the native state. |
|-------------------|--|
| Applications: | ELISA, IHC |
| Name of antibody: | TBCC |
| Immunogen: | Fusion protein of human TBCC |
| Full name: | tubulin folding cofactor C |
| Synonyms: | CFC |



| SwissProt: | Q15814 |
|------------------------------------|-----------------------|
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human prostate cancer |
| IHC Recommend dilution: | 100-200 |

