

兔抗 PTPN6(Ab-564) 多克隆抗体

中文名称: 兔抗 PTPN6(Ab-564) 多克隆抗体

英文名称: Anti-PTPN6(Ab-564) rabbit polyclonal antibody

别 名: MSK1; RLPK; MSPK1

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: PTPN6(Ab-564)

反应种属: Human

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

n of the transcription factors CREB1 and ATF1 a nd for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contribute s to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes. Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C

Serine/threonine-protein kinase that is required f or the mitogen or stress-induced phosphorylatio

irradiation, epidermal growth factor (EGF) and an



www.shjning.com

| Synonyms: | 5 MSK1; RLPK; MSPK1 |
|---------------------------------|--|
| Full name: | ribosomal protein S6 kinase, 90kDa, polypeptide |
| Immunogen: | Synthesized peptide derived from C-terminal of human SHP-1. |
| мате от аптіроду: | |
| | |
| Applications: Name of antibody: | isomycin. Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytopl asm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression. In skeletal myobla sts is required for phosphorylation of RELA at 'S er-276' during oxidative stress. In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and regulation of its transcriptional potential. Phosphorylates ETV1/ER81 at 'Ser-191' and 'Ser-216', and thereby regulates its ability to stimulate transcription, which may be important during development and breast tum or formation. Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A. Phosp horylates 'Ser-10' of histone H3 in response to mitogenics, stress stimuli and EGF, which results in the transcriptional activation of several immed iate early genes, including proto-oncogenes c-fo s/FOS and c-jun/JUN. May also phosphorylate 'S er-28' of histone H3. Mediates the mitogen- and stress-induced phosphorylation of high mobility group protein 1 (HMGN1/HMG14). In lipopolysac charide-stimulated primary macrophages, acts do wnstream of the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines. Functions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inflammatory cytokine interleukin 10 (IL10), via CRE B1 and ATF1 transcription factors. Plays a role in neuronal cell death by mediating the downstream effects of excitotoxic injury. WB PTPN6(Ab-564) |



www.shjning.com

| SwissProt: | O75582 |
|--------------------------|---------------------|
| WB Predicted band size: | 90 kDa |
| WB Positive control: | Jurkat cells lysate |
| WB Recommended dilution: | 500-3000 |

