

兔抗 TCIM 多克隆抗体

- 中文名称: 兔抗 TCIM 多克隆抗体
- 英文名称: Anti-TCIM rabbit polyclonal antibody
- 别 名: transcriptional and immune response regulator; TC1
- 相关类别: 一抗
- 储 存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗 原: TCIM
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a small, monomeric, predominantly unstructured protein that functions as a positive regulat or of the Wnt/beta-catenin signaling pathway. This prot ein interacts with a repressor of beta-catenin mediated transcription at nuclear speckles. It is thought to comp etitively block interactions of the repressor with beta-ca tenin, resulting in up-regulation of beta-catenin target genes. The encoded protein may also play a role in th e NF-kappaB and ERK1/2 signaling pathways. Expressio n of this gene may play a role in the proliferation of s everal types of cancer including thyroid cancer, breast cancer and hematological malignancies. [provided by R efSeq, Nov 2011]
Applications:	ELISA, IHC
Name of antibody:	TCIM



Immunogen:	Fusion protein of human TCIM
Full name:	transcriptional and immune response regulator
Synonyms:	TC1; TC-1; C8orf4
SwissProt:	Q9NR00
IHC positive control:	Human gastric cancer and Human thyroid cancer
IHC Recommend dilution:	40-200





