

兔抗 STOML2 多克隆抗体

中文名称：兔抗 STOML2 多克隆抗体

英文名称：Anti-STOML2 rabbit polyclonal antibody

别名：SLP-2; HSPC108

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：STOML2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Stomatin-like protein 2 (STOML2), SLP-2 or EPB72-like 2, is a 356 amino acid member of the mec-2 family of proteins. Expressed ubiquitously at low levels, STOML2 is highly expressed in heart, liver and pancreas. STOML2 is localized to the cytoplasm with some distribution on the membrane. STOML2 was first identified as an overexpressed protein in human endometrial adenocarcinoma. Changes in cell growth in samples with different levels of STOML2 indicate that STOML2 could play a role in endometrial tumorigenesis. STOML2 is also thought to play a role in regulating ion channel conductances or the organization of sphingolipid and cholesterol-rich lipid rafts.
Applications:	ELISA, WB, IHC
Name of antibody:	STOML2

Immunogen:	Fusion protein of human STOML2
Full name:	stomatin like 2
Synonyms:	SLP-2; HSPC108
SwissProt:	Q9UJZ1
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human prostate cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	39 kDa
WB Positive control:	Mouse brain tissue ,Rat heart tissue ,Rat brain tissue ly es
WB Recommended dilution:	500-2000



