

兔抗 SHC1 (Phospho-Tyr349)多克隆抗体

中文名称: 兔抗 SHC1 (Phospho-Tyr349)多克隆抗体

英文名称: Anti-SHC1 (Phospho-Tyr349) rabbit polyclonal antibody

别 名: SHC; SHCA

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: SHC1 (Phospho-Tyr349)

反应种属: Human, Mouse

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

Signaling adapter that couples activated growth fa ctor receptors to signaling pathway. Isoform p46Sh c and isoform p52Shc, once phosphorylated, coupl e activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are imp licated in the cytoplasmic propagation of mitogeni c signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling casc ade in various non-neuronal systems. Isoform p66S hc does not mediate Ras activation, but is involve d in signal transduction pathways that regulate the cellular response to oxidative stress and life span. I soform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation



www.shjning.com

| | of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has b een correlated with life span |
|--------------------------|---|
| Applications: | WB, IF |
| Name of antibody: | SHC1 (Phospho-Tyr349) |
| Immunogen: | Synthetic peptide of human SHC1 (Phospho-Tyr349) |
| Full name: | SHC (Src homology 2 domain containing) transfor ming protein 1 (Phospho-Tyr349) |
| Synonyms: | SHC; SHCA |
| SwissProt: | P29353 |
| WB Predicted band size: | 63 kDa |
| WB Positive control: | HepG2 cells treated with EGF |
| WB Recommended dilution: | 500-1000 |
| IF Positive control: | Hela cells |
| IF Recommended dilution | 100-200 |





