

兔抗 SIGLEC7 多克隆抗体

中文名称：兔抗 SIGLEC7 多克隆抗体

英文名称：Anti-SIGLEC7 rabbit polyclonal antibody

别名：p75; QA79; AIRM1; CD328; CDw328; D-siglec; SIGLEC-7; SIGLECP2; SIGLEC19P; p75/AIRM1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SIGLEC7

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Siglec-7, which is highly expressed in monocytes and resident blood cells but not in parenchymatous cells, mediates inhibition of natural killer cell cytotoxicity. Due to alternative splicing events, two isoforms exist for Siglec-12, namely SLG-L (the long isoform) and SLG-S (the shorter isoform). These isoforms are differentially expressed with the longer isoform predominantly found in small intestine, spleen and bone marrow, and the shorter isoform predominantly found in small intestine, spleen and adrenal gland.
Applications:	ELISA, IHC
Name of antibody:	SIGLEC7
Immunogen:	Synthetic peptide of human SIGLEC7

Full name:	sialic acid binding Ig-like lectin 7
Synonyms :	p75; QA79; AIRM1; CD328; CDw328; D-siglec; SIGLEC-7; SIGLECP2; SIGLEC19P; p75/AIRM1
SwissProt:	Q9Y286
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human thyroid cancer and Human brain
IHC Recommend dilution:	25-100

