

兔抗 MED15 多克隆抗体

中文名称: 兔抗 MED15 多克隆抗体

英文名称: Anti-MED15 rabbit polyclonal antibody

别名: TIG1; CAG7A; CTG7A; PCQAP; TIG-1; TNRC7; ARC105

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: MED15

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

| Background: | The protein encoded by this gene is a subunit of the m ultiprotein complexes PC2 and ARC/DRIP and may functi on as a transcriptional coactivator in RNA polymerase II transcription. This gene contains stretches of trinucleotide repeats and is located in the chromosome 22 region whi ch is deleted in DiGeorge syndrome. Alternative splicing results in multiple transcript variants. |
|-------------------|---|
| Applications: | ELISA, IHC |
| Name of antibody: | MED15 |
| Immunogen: | Synthetic peptide of human MED15 |
| Full name: | mediator complex subunit 15 |
| Synonyms: | TIG1; CAG7A; CTG7A; PCQAP; TIG-1; TNRC7; ARC105 |
| SwissProt: | Q96RN5 |



| ELISA Recommended dilution: | 2000-5000 |
|-----------------------------|----------------------|
| IHC positive control: | Human ovarian cancer |
| IHC Recommend dilution: | 25-100 |

