

## 兔抗 PDCD10 多克隆抗体

中文名称：兔抗 PDCD10 多克隆抗体

英文名称：Anti-PDCD10 rabbit polyclonal antibody

别名：CCM3, TFAR15

抗原：PDCD10

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

|                          |   |
|--------------------------|---|
| <b>Background:</b>       | This gene encodes an evolutionarily conserved protein associated with cell apoptosis. The protein interacts with the serine/threonine protein kinase MST4 to modulate the extracellular signal-regulated kinase (ERK) pathway. It also interacts with and is phosphorylated by serine/threonine kinase 25, and is thought to function in a signaling pathway essential for vascular development. Mutations in this gene are one cause of cerebral cavernous malformations, which are vascular malformations that cause seizures and cerebral hemorrhages. Multiple alternatively spliced variants, encoding the same protein, have been identified. |
| <b>Applications:</b>     | ELISA, WB, IHC  |
| <b>Name of antibody:</b> | PDCD10  |

|                                    |                                     |
|------------------------------------|-------------------------------------|
| <b>Immunogen:</b>                  | Fusion protein of human PDCD10      |
| <b>Full name:</b>                  | programmed cell death 10            |
| <b>Synonyms :</b>                  | CCM3, TFAR15                        |
| <b>SwissProt:</b>                  | Q9BUL8                              |
| <b>ELISA Recommended dilution:</b> | 1000-5000                           |
| <b>IHC positive control:</b>       | Human tonsil and human renal cancer |
| <b>IHC Recommend dilution:</b>     | 25-100                              |
| <b>WB Predicted band size:</b>     | 25 kDa                              |
| <b>WB Positive control:</b>        | Hela, Jurkat and 231 cells          |
| <b>WB Recommended dilution:</b>    | 200-1000                            |



