

兔抗 PDIA5 多克隆抗体

中文名称: 兔抗 PDIA5 多克隆抗体

英文名称: Anti-PDIA5 rabbit polyclonal antibody

别 名: protein disulfide isomerase family A member 5; PDIR

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: PDIA5

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange re actions. The encoded protein has an N-terminal ER-signa I sequence, three catalytically active thioredoxin (TRX) do mains, a TRX-like domain, and a C-terminal ER-retention sequence. The N-terminal TRX-like domain is the primary binding site for the major ER chaperone calreticulin and possibly other proteins and substrates as well. Alternative splicing results in multiple protein- and non-protein-coding transcript variants.
Applications:	ELISA, WB, IHC
Name of antibody:	PDIA5



www.shjning.com

Immunogen:	Fusion protein of human PDIA5
Full name:	protein disulfide isomerase family A member 5
Synonyms:	PDIR
SwissProt:	Q14554
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	30-150
WB Predicted band size:	60 kDa
WB Positive control:	Human placenta tissue lysate
WB Recommended dilution:	1000-5000







